

JOINT COLLOQUIA

27 April 1967

THE 26-MONTH OSCILLATION OF METEOROLOGICAL ELEMENTS
IN THE STRATOSPHERE

By Jennifer Hopwood (Bureau of Meteorology, Melbourne)

This now well documented phenomenon (Reed, 1964) concerns zonal wind, temperature and ozone, but the speaker dealt mainly with the wind. The subject was introduced with an outline of the prominent equatorial oscillation and its mode of poleward attenuation in the northern hemisphere. To describe a dissimilar behaviour in the southern hemisphere that had emerged from recent studies (Laby, Sparrow and Unthank, 1964; Sparrow and Unthank, 1964; Berson, 1966; Rofe, 1966) Mrs. Hopwood then presented a harmonic analysis of seven years' observations from the Australian region. Whereas there was marked attenuation southward from Singapore through Lae and Darwin on to Alice Springs (where the cycle is practically non-existent through almost the entire stratosphere), it was again detectable as a modulation of the strong annual cycle at middle latitude stations such as Laverton and Hobart.

Analysis of variance indicated that the middle latitude phenomenon was not a statistical accident. A noteworthy feature of the oscillation in southern middle latitudes was the absence of downward phase propagation characteristic of the tropical stratosphere (approx. 1 km/month).

Discussing the mechanics of vertical and horizontal momentum transfers associated with the equatorial oscillation (Tucker, 1964), the speaker gave results of numerical solutions which were based on slightly idealized profiles of the mean zonal and vertical velocities. An assumption underlying the model (Tucker, 1964) was the time independence of the momentum transfer coefficient. In current work the treatment is being extended to incorporate a varying coefficient. It was hoped that the study of the middle latitude cycle and its relation to the tropical oscillation would lead to an increased understanding of the whole phenomenon.

In the general discussion Mr. Rider (Aeronautical Research Laboratory) raised the question whether and, if so, how, in the model, the vertical motion was coupled with the oscillatory nature of both zonal momentum and lateral eddy-transfer of momentum. Mrs. Hopwood explained that there was no mathematical inconsistency in the model, and Dr. Tucker (Bureau of Met.) amplified this by pointing out that vertical transport of momentum was allowed to vary on account of time dependent vertical wind shear, even though the transfer coefficient was assumed constant.

Dr. Berson (C.S.I.R.O.) referred to the speaker's mention of variable ozone heating as one of the possible energy sources for the oscillation and questioned whether such would still be consistent with the basic feature of the model in which the oscillation was maintained by lateral transfer of momentum. Dr. Tucker clarified some points concerning this important problem of momentum generation and this led to a more general discussion of related aspects of the oscillation.

It was pointed out by Dr. Loewe that the oscillation phenomenon was indeed world-wide and has its counterpart in quasi-biennial variations of meteorological surface elements detected at some places over 50 years ago. Other questions raised were the present decline of amplitude outside the equatorial belt as opposed to the equatorial regime where it was apparently maintained; the relationship with sunspot cycle; and connections with terrestrial causes of radiation changes, especially due to variation of cloudiness in the tropics possessing a semi-annual cycle which approximates a sub-harmonic of the 26-month cycle.

REFERENCES

- | | | |
|--|------|---------------------------------------|
| Reed, R.J. | 1964 | Q.J.R. Met. Soc. <u>90</u> , p. 441 |
| Laby, J.E., Sparrow, J.G.
& Unthank, E.L. | 1964 | J. Atmos. Sci. <u>21</u> , p. 249-259 |

Sparrow, J.G., & Unthank, E.L.	1964	J. Atmos. Sci. <u>21</u> , p. 592.
Rofe, B.	1966	W.R.E. Technical Note Pad 115.
Berson, F.A.	1966	Nature, 5042, p. 1243.
Tucker, G.B.	1964	Q.J.R. Met. Soc., <u>90</u> , p. 405.

F.A.B.

25 May, 1967

FINITE DIFFERENCE OPERATORS AND THEIR USE IN NUMERICAL FORECASTING

by D. Jenssen

Dr. Ditmar Jenssen of the Meteorology Department, University of Melbourne, opened his talk with a lengthy preamble setting the background for the work to be discussed. The role of finite difference approximations to various simple partial derivatives was examined very generally. Examples were selected from the fields of Numerical forecasting, integration, iterative solution of partial D.E.'s, estimation of a numerical value of complex differential functions, and finally two-dimensional interpolation.

The derivation of the difference analogues was then briefly outlined. Using a one-dimensional Taylor's series it was shown how standard operators were developed: the theory was then extended to two-dimensions, and the importance of cross terms such as $\partial^2/\partial x \partial y$ indicated. Analogues to ∇^2 and $\partial/\partial x$ were discussed in relative detail. The results of the testing of these operators on single components of a double Fourier field followed. Slides were shown giving the dependency of the analogue accuracy on the wavelength scale of the field upon which they act.

The formulation of a mathematical simulation of a 500 mb chart was then treated, and its features discussed. The results of applying the analogues to such 'ideal' meteorological data were shown and the best difference operators for this typical flow pattern determined.

A brief treatment of the use of the various finite difference approximations to the Laplacian to the iterative solution of Poisson's equation was then undertaken. It was shown that the accuracy of the solution was determined not primarily by the number of iterations (provided this is sufficiently large) but by the accuracy of the analogue employed. The speed of solution was also discussed, and by making a compromise between speed and accuracy the most efficient analogue for utilisation in this iterative scheme was chosen. The talk concluded with an outline of further planned work. This included the possible development of three- and four-dimensional operators, and a detailed examination of the iterative solution of equations other than Poisson's, particularly those involving the bi-harmonic operator.

During the question period, it was suggested by Mr. Wallington that a profitable further line of investigation would be the topic of 'compatibility': that is, would the analogues, when substituted into a differential equation, change the nature of that equation to such an extent that the results obtained were virtually useless?

Dr. Radok suggested that it may be more expedient to specify meteorological fields by two-dimensional Fourier analysis, rather than by a direct mathematical simulation such as the one discussed in the talk.

The usefulness of high-accuracy analogues in numerical forecasting, when the initial data could have a relatively high error, was discussed by Mr. Budd, other participants and the speaker. It was decided that the better the difference operator, the less likely it would be that the initial errors would amplify, and therefore the better the forecast would be.

Dr. Berson referred to the slide giving analogue accuracy as a function of wavelength, and pointed out that the wavelength of, say, the 500 mb field would not be the same in both the x and y directions. Dr. Jenssen agreed this would be so, but stated that the results obtained showed that the shorter wavelength was the dominant one in fixing the accuracy of the analogue, and it was therefore legitimate to speak of analogue accuracy as a function of a single wavelength.