

**THE SOUTH PACIFIC SEA LEVEL & CLIMATE
MONITORING PROJECT**

MONTHLY DATA REPORT

NO. 160

OCTOBER 2008



Australian Government

Bureau of Meteorology

This project is sponsored by the Australian Agency for International Development (AusAID), and is managed by the Bureau of Meteorology with its National Tidal Centre (NTC) providing key technical support.



Australian Government

Bureau of Meteorology

National Tidal Centre Bureau of Meteorology Australia

GPO Box 421
Kent Town SA 5071
Australia

Tel: (+618) 8366 2730
Fax: (+618) 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Quality Certification:

I authorise the issue of this South Pacific Sea Level and Climate Monitoring Project Monthly Data Report for October 2008 in accordance with National Tidal Centre Quality Assurance procedures.

William Mitchell
Manager - National Tidal Centre

South Pacific Sea Level and Climate Monitoring Project

Monthly Data Report

October 2008

EXECUTIVE SUMMARY

This summary, and the overview that follows, are intended to provide a synopsis of the Monthly Data Report and of the trends observed over the life of the project to date.

October 2008

- The SEAFRAME network continued to collect high quality sea level and associated meteorological information for monitoring climate variability and climate change.
- The monthly mean sea level at Vanuatu was the highest on record, eclipsing the previous record set in December 2001 by 3mm.
- The sea level observations at FSM and PNG during October 2008 showed instances of natural harbour resonance.
- Slightly higher than normal sea levels and water temperatures continued to be observed at the southern stations of Vanuatu, Fiji, Tonga and Cook Islands. Sea levels remained lower than normal at the equatorial stations Kiribati and Nauru but water temperatures there have returned to near normal following an extended period of cooler than normal conditions.
- Climate conditions across the equatorial Pacific remain neutral despite various La Niña characteristics such as positive Southern Oscillation Index values and stronger than normal Trade Winds being observed.
- The majority of international climate models predict that neutral climate conditions will persist into early 2009.

Short-Term Trends

It is important to stress that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Observed trends in sea level include natural variability, for example, events such as El Niño and effects due to many other atmospheric, oceanographic and geological processes. Longer-term data sets for all

stations are required in order to separate the effects of the different signals. ***Please exercise caution in interpreting the short-term trends in the table below*** – they will almost certainly change over the coming years as the data set increases in length. Figure 13 later in this report provides the “time history” of the short-term trend at all project locations.

Recent short-term sea level trends in the project area based upon SEAFRAME data through October, 2008				
Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cook Is	21°12'17.1"S / 159°47'5.2"W	Feb 1993	+5.2	+0.1
Tonga	21°8'12.5"S / 175°10'50.5"W	Jan 1993	+9.2	+0.1
Fiji	17°36'17.7"S / 177°26'17.7"E	Oct 1992	+4.5	+0.1
Vanuatu	17°45'19.2"S / 168°18'27.7"E	Jan 1993	+4.6	+0.2
Samoa	13°49'36.4"S / 171°45'40.7"W	Feb 1993	+6.1	-0.1
Tuvalu	8°30'8.9"S / 179°11'42.6"E	Mar 1993	+5.8	-0.1
Kiribati	1°21'54.2"N / 172°55'58.8"E	Dec 1992	+3.7	-0.2
Nauru	0°31'45.9"S / 166°54'36.2"E	Jul 1993	+4.4	-0.2
Solomon Is.	9°25'44.1"S / 159°57'19.3"E	Jul 1994	+7.9	0.0
PNG	2°2'31.5"S / 147°22'25.6"E	Sep 1994	+7.8	0.0
FSM	6°58'49.9"N / 158°12'0.8"E	Dec 2001	+21.6	-0.3
Marshall Is.	7°6'21.7"N / 171°22'22.1"E	May 1993	+4.0	+0.1

INTRODUCTION

Welcome to the October 2008 Monthly Data Report for the South Pacific Sea Level and Climate Monitoring Project (SPSLCMP). The report details the month by month operation of the SEAFRAME monitoring stations in the Pacific, including operational problems with the network or with satellite communications, the occurrence of abnormal sea level or climate events, interpretation of sea level fluctuations in the context of El Niño and the emergence of trends in the data.

The SPSLCMP was developed as an Australian response to concerns raised by the member countries of the South Pacific Forum over the potential impacts of global warming on climate and sea levels in the Pacific. Support was provided for the installation of SEAFRAME monitoring stations across the South Pacific Forum region.

SEAFRAME gauges not only measure sea level by two independent means, but also observe a number of “ancillary” variables - air and water temperatures, wind speed, wind direction and atmospheric pressure. There is an associated programme of levelling to first order, to determine shifts in the vertical of the sea level sensors due to local land movement. Continuous Global Positioning System (CGPS) measurements

are now also being made to determine the vertical movement of the land with respect to the International Terrestrial Reference Frame.

The AusAID funded project has, as its principal objective *‘the provision of an accurate long term record of sea level in the South Pacific for partner countries and the international scientific community, that enables them to respond to and manage related impacts’*.

The project’s monitoring network consists of 12 SEAFRAME stations, providing a wide coverage across the Southwest Pacific basin. All of these stations (see Figure A), with the exception of the Pohnpei (FSM) gauge, which was established in December 2001, have been operational since October 1994.

The monthly data report, one of a range of information products produced by the project, is the primary form of SPSLCMP data dissemination. Its content is designed to provide up-to-date access to the project’s data products.




Figure A: *South Pacific Sea Level and Climate Monitoring Stations*

OCTOBER CLIMATOLOGY

Neutral El Niño / Southern Oscillation climate conditions continued to be observed across the equatorial Pacific during October. Sea surface temperatures across the equatorial Pacific were near normal, while subsurface temperatures in the eastern equatorial Pacific remained cooler than normal. Climate model forecasts suggest neutral conditions will likely persist until early next year.

The Southern Oscillation Index (SOI) remains positive, with the October monthly value of +13 following the September value of +14 (**Figure B**). Despite positive SOI values the potential for La Niña conditions to develop this year is considered low.

Maps of the sea surface temperature anomalies (**Figure C, Figure E**) show near average conditions prevailed across most of the equatorial Pacific during October. Surface temperatures in the far eastern and far western equatorial Pacific were slightly warmer than normal. A band of warmer than normal surface temperatures continued to be observed across the southwest Pacific.

Subsurface temperatures remained cooler than normal across the central and eastern equatorial Pacific during October, although some subsurface warming occurred across the central equatorial Pacific (**Figure D**).

During El Niño (warm-episode) conditions there is a sustained weakening of the Trade Winds across much of the equatorial Pacific and an increase in cloudiness in the central Pacific particularly near the dateline. During La Niña (cold-episode) conditions there is a reversal of this situation, with stronger Trade Winds and a decrease in cloudiness in the central Pacific. Trade Winds during October continued to be stronger than normal across the central and western equatorial Pacific (**Figure E**). Trade Winds across the eastern equatorial Pacific were near to slightly higher than normal. Cloudiness in the equatorial Pacific near the dateline continued to be slightly below normal during October in keeping with slightly cooler than normal sea surface temperatures in that region.

The consensus among six international computer models is that neutral climate conditions will persist across the equatorial Pacific into early 2009.

The preceding description of the climatology of the Pacific region, and Figures B, C and D are based on information sourced from the National Climate Centre of the Australian Bureau of Meteorology at <http://www.bom.gov.au/climate/>. Figure E was generated from the Tropical Atmosphere Ocean project website courtesy of PMEL, NOAA at <http://www.pmel.noaa.gov/tao/>.




Figure B: The five-month weighted mean and individual monthly means of the Southern Oscillation Index (SOI). The SOI is ten times the monthly anomaly of the difference in mean sea level pressure between Tahiti and Darwin, divided by the long-term standard deviation of that difference for the relevant month.




Figure C: Sea surface temperature anomaly ($^{\circ}\text{C}$) for October 2008.




Figure D: Equatorial depth-longitude section of ocean temperature anomalies for July 2008 through to October 2008. Contour interval is 0.5°C .




Figure E: Monthly mean wind vectors (top) and anomalies (bottom) for October 2008. The colour-shaded contours represent the monthly mean sea surface temperatures (top) and anomalies (bottom).

OCTOBER SEAFRAME DATA

Monthly Sea Level and Environmental Data (Figures 1-10)

The **observed sea levels (Figure 1)** are dominated by the daily oscillations of the tide. In most cases, the tide rises and falls twice per day (semi-diurnal), but at PNG and the Solomon Islands the tide tends to have a single high and low per day (diurnal). The greatest variations are called spring tides and tend to occur close to the full and new moon. There was a full moon on the 14th of October and a new moon on the 28th of October UTC.

Gaps in the data are the result of instrumental errors or data retrieval problems and are discussed under **Instrument Performance**.

The **residuals (Figure 2)** are the differences between the observed sea levels and the tidal predictions. They highlight the non-tidal sea level fluctuations such as those due to the short-term effects of weather or tsunamis. Residual sea level fluctuations may also be amplified or sustained by the shape of the harbour in which the gauge is located. Persistent sloshing of water within a bay or harbour, for example, is known as a seiche. Seiches are often recorded at PNG when the wind suddenly changes strength or direction, which happened on a daily basis around the 7th-11th October. The residual sea levels at FSM occasionally become prominent around the time of neap tides, and several good examples of this were observed during October.

The sea level residuals at all stations, to some degree, exhibit semi-diurnal or diurnal fluctuations, which last a few days or weeks and then disappear. If these peaks were to persist, rather than appear as occasional ‘transients’, then the tidal analysis would be able to account for them, and the end result would be virtual eradication from the residuals.

The **barometrically corrected residuals (Figure 3)** have had the effect of atmospheric pressure fluctuations removed from the sea level residuals of Figure 2. The rule of thumb for the ‘inverse barometer effect’ is that a 1-hPa fall in the barometer, if sustained over a day or more, produces a 1 cm rise in the local sea level (within the area beneath the low pressure system).

The **winds, temperatures and barometric pressures** are plotted in **Figures 4 to 9**. The short lines in **Figure 5** follow the meteorological convention, that is, they point in the direction the wind is coming *from*. For example, the winds at Vanuatu prevailed from the northeast for most of the month. Periods of strong winds were observed at Fiji in the first half of October, including gusts to 25 m/s (49 knots or 90 km/hr) on the 10th of October (**Figure 6**).

Air and water temperatures (**Figures 7 and 8**) are plotted using the same vertical scale for the purpose of comparison. The air temperatures are seen to fluctuate over a much wider range than the water temperatures. At some sites (e.g. FSM) the water temperature shows almost no variation, although the air temperature varies by several degrees between night and day. At Nauru a twice-daily fluctuation in water temperature is evident and is related to the tide, as it is usually more pronounced during the larger spring tides.

Barometric pressures (**Figure 9**) tend to fluctuate by around 3 hPa twice-daily at all stations as a result of atmospheric tides, which are largest in the tropical regions and reduce to near zero toward the poles. The longer-term barometric pressure fluctuations that occur over periods of days to weeks are due to passing weather systems. These fluctuations tend to be larger at sites further away from the equator such as Cook Islands and Tonga.

The **meteorological data** are put into perspective by **Figure 10**. In this figure, if an open circle falls above (below) a solid dot, a new maximum (minimum) for the particular month has been set. *The data sets only include South Pacific Sea Level and Climate Monitoring Project data, which have been collected since October 1992 when the first station was installed (Fiji). The data from FSM has only been collected since December 2001.* A new maximum October air temperature of 32.1°C was recorded at Marshall Islands and a new maximum October water temperature of 27.7°C was observed at Cook Islands.

Mean Sea Level and Anomalies (Figures 11-13)

Figure 11 shows the **monthly mean sea levels**, which are simple arithmetic averages of the sea levels, relative to an arbitrary zero. The figure shows that Tuvalu, for example, normally experiences an annual cycle of about 0.2 metres, reaching a peak around February or March. One effect of the El Niño of 1997/1998 was to disrupt the annual sea level cycle at many of the SEAFRAME stations.

The monthly mean sea level at Vanuatu for October 2008 of 1.017m is the highest on record. The previous highest monthly mean sea level of 1.014m was observed in December 2001.

Figure 12 shows the monthly mean **sea level anomalies**, or departures from normal conditions after tides, annual and semi-annual seasonal cycles and the sea level trend have been removed. The annual cycle at Tuvalu (which has the largest consistent annual cycle) is quite notable in **Figure 11** but less apparent in **Figure 12**. By removing the seasonal cycles, the anomalies help to bring out irregular features, such as lower than normal sea levels across the region during the 1997/98 El Niño.

In October 2008 sea levels 5-10cm higher than normal continued to be observed at Vanuatu, Fiji, Tonga and Cook Islands. The equatorial stations of Nauru and Kiribati continued to experience sea levels at least 5cm below normal. Lower than normal sea levels were also observed at Tuvalu and Samoa. Sea levels at the remaining stations were generally within 5cm of what is normally observed for this time of the year.

Sea Level Trends

The **short-term sea level trends** at individual stations as at October 2008 are shown in the following table. Sea level trends are updated every month by allowing for a linear trend term in the tidal analysis of all the data available at individual stations. *Please exercise caution in interpreting the trends* – they will continue to change over the coming years as the data sets increase in length. The evolution of the monthly trend values (in mm per year) at each station from one year after installation to present is depicted in **Figure 13**. This figure illustrates that as the sea level record becomes longer, the relative sea level trend estimates become more stable and reliable. The

reason for this is that the trends from short sea level records are affected by the natural sea level variability occurring on inter-annual, El Niño and decadal timescales due to atmospheric, oceanographic and geological processes. Longer-term data sets for all stations are required in order for the underlying trend to emerge from these short-term variations. Further details are available from the *National Tidal Centre (NTC)*, *Australian Bureau of Meteorology*.

Recent short-term sea level trends in the project area based upon SEAFRAME data through October, 2008				
Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cook Is	21°12'17.1"S / 159°47'5.2"W	Feb 1993	+5.2	+0.1
Tonga	21°8'12.5"S / 175°10'50.5"W	Jan 1993	+9.2	+0.1
Fiji	17°36'17.7"S / 177°26'17.7"E	Oct 1992	+4.5	+0.1
Vanuatu	17°45'19.2"S / 168°18'27.7"E	Jan 1993	+4.6	+0.2
Samoa	13°49'36.4"S / 171°45'40.7"W	Feb 1993	+6.1	-0.1
Tuvalu	8°30'8.9"S / 179°11'42.6"E	Mar 1993	+5.8	-0.1
Kiribati	1°21'54.2"N / 172°55'58.8"E	Dec 1992	+3.7	-0.2
Nauru	0°31'45.9"S / 166°54'36.2"E	Jul 1993	+4.4	-0.2
Solomon Is.	9°25'44.1"S / 159°57'19.3"E	Jul 1994	+7.9	0.0
PNG	2°2'31.5"S / 147°22'25.6"E	Sep 1994	+7.8	0.0
FSM	6°58'49.9"N / 158°12'0.8"E	Dec 2001	+21.6	-0.3
Marshall Is.	7°6'21.7"N / 171°22'22.1"E	May 1993	+4.0	+0.1

Barometric Pressure, Water Temperature and Air Temperature Anomalies

The anomalies of barometric pressure, water and air temperature (**Figures 14 to 16**) are determined in the same manner as the sea level anomalies (**Figure 12**), except the trend is not calculated.

The **barometric pressure anomalies** (**Figure 14**) show substantially higher than normal barometric pressures were observed at SEAFRAME stations during the 1997-1998 El Niño. In October 2008 barometric pressures were near to slightly higher than normal at most stations, with Cook Islands showing the largest anomaly of 2hPa.

The **water temperature anomalies** (**Figure 15**) show a return to near normal water temperatures was observed at the equatorial stations of Kiribati and Nauru during October 2008 following an extended period of cooler than normal conditions. It is not unusual for observations near the equator to differ from the other stations as a result of large-scale waves that propagate along the equatorial waveguide. Warmer than normal water temperatures were observed at the southern stations of Vanuatu, Fiji, Tonga and Cook Islands and accompany the higher than normal sea levels at those stations.

Warmer than normal conditions were also observed at PNG and Solomon Islands, while cooler than normal water temperatures were observed at Marshall Islands, FSM, Tuvalu and Samoa.

The **air temperature anomalies (Figure 16)** during October 2008 show warmer than normal conditions were observed at Marshall Islands, PNG, Vanuatu, Fiji, Tonga and Cook Islands. Conditions at FSM, Solomon Islands, Kiribati, and Samoa were generally near to what is normally observed at this time of the year. At Nauru and Tuvalu no air temperature data were received. Over the duration of the record the air temperature anomalies generally (although not always) follow the water temperature anomalies, which is an indication of the large influence the ocean has upon the climate of the Pacific Islands.

Instrument Performance

In **Figure 17**, which shows **sea level data return**, colour is used to distinguish five-year project phases. The number of missing days is noted in gaps in the bars. Very good sea level data return was achieved in October 2008.


The SEAFRAME stations at Fiji and Tuvalu were serviced in October 2008 as part of the scheduled calibration and maintenance program, resulting in some small data gaps at those stations. The air temperature sensors at Nauru and Tuvalu continue to experience problems and erroneous data were removed from the record. The air temperature, water temperature and barometric pressure observations at Marshall Island contained numerous isolated errors or 'spikes' that were removed from the record. Similar treatment was necessary for the water temperature observations at Kiribati and Tonga. At Nauru erroneous sea level and water temperature observations were replaced with observations from backup sensors.

SEAFRAME STATIONS

SEAFRAME stations employ a SUTRON programmable data logger, water level gauges and other sensors. The data logger and associated electronics are normally housed in fibreglass huts. A sketch of a typical station is shown in the following figure. Water level sensors include:


- (1) Primary water level using a Bartex 'AQUATRAK' acoustic-in-air sensor, and
- (2) Secondary water level (or backup) using a Druck pressure transducer mounted close to the seabed.

The primary and backup water level sensors provide water level values, which are averaged over three minutes and are logged every six minutes. The data logger has the memory capacity to store approximately one month of data. The meteorological sensors are logged to the SUTRON data logger on an hourly basis.



Web Hits

The following chart shows the number of times the Pacific pages on the *NTC* web site have been visited, by month since January 2005. Note that the web statistics for February 2005 are not available due to technical difficulties.



The *Monthly Data Report* is prepared by *NTC* for *AusAID*.

NTC would appreciate feedback from readers on the content and presentation of the *Monthly Data Report*.

Please spare a few moments to let us know your constructive opinion.

Further communication on the *Monthly Data Report* may be made to *NTC*. Anyone interested in a more detailed account of the project should contact:

National Tidal Centre
Bureau of Meteorology
PO Box 421
Kent Town SA 5067
Tel: (+618) (08) 8366 2600
Fax: (+618) (08) 8366 2693
Website: <http://www.bom.gov.au/oceanography/tides.shtml>

Or visit the project website at <http://www.bom.gov.au/pacificsealevel>

Please refer to: <http://www.bom.gov.au/oceanography/projects/spslcmp/spslcmp.shtml> for details.

Please also note the following:

While care has been taken in the collection, analysis, and compilation of the data, it is supplied on the condition that neither the *Commonwealth of Australia* nor *NTC* shall be liable for any loss or injury whatsoever arising from the use of the data. Copyright for material contained in this document is held by the *Commonwealth of Australia*.

Individuals and organisations are advised that quality controlled six-minute or hourly data from these stations are available on request from *NTC*. Some handling fees may be charged. For commercial agencies requesting data, some additional costs may be levied.

Figure 1

OCTOBER 2008

SIX MINUTE WATER LEVEL OBSERVATIONS (m)




Figure 2

OCTOBER 2008
SIX MINUTE RESIDUAL WATER LEVELS (m)




Figure 3




Figure 4

OCTOBER 2008
HOURLY WIND SPEEDS (m/s)




Figure 5
OCTOBER 2008
HOURLY INCIDENT WINDS (m/s, deg True)

— 10 m/s




Figure 6
OCTOBER 2008
HOURLY MAXIMUM WIND GUSTS (m/s)




Figure 7

OCTOBER 2008
HOURLY AIR TEMPERATURES (°C)




Figure 8

OCTOBER 2008
HOURLY WATER TEMPERATURES (°C)




Figure 9
OCTOBER 2008
HOURLY ATMOSPHERIC PRESSURE (hPa)







Figure 10

Comparison of October 2008 Max, Min & Mean with Long Term October Values



Stations

- 1 - Marshall Islands
- 2 - Federated States of Micronesia
- 3 - Papua New Guinea
- 4 - Solomon Islands
- 5 - Kiribati
- 6 - Nauru
- 7 - Tuvalu
- 8 - Samoa
- 9 - Vanuatu
- 10 - Fiji
- 11 - Tonga
- 12 - Cook Islands

- October 2008 Maximum
- October 2008 Mean
- October 2008 Minimum
- Long Term October Maximum
- Long Term October Mean
- Long Term October Minimum

Figure 11

MONTHLY MEAN SEA LEVELS TO OCTOBER 2008 (m)

The zero line represents an arbitrary fixed offset from the zero of the tide gauge.




Figure 12

SEA LEVEL ANOMALIES THROUGH OCTOBER 2008 (m)




Figure 13

SEA LEVEL TRENDS THROUGH OCTOBER 2008 (mm/year)





Figure 14

BAROMETRIC PRESSURE ANOMALIES THROUGH OCTOBER 2008 (hPa)



1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008



Figure 16
**AIR TEMPERATURE ANOMALIES
 THROUGH OCTOBER 2008 (°C)**




Figure 17

SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF GAP ARE INDICATED
GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE

* Patchy record

