THE SOUTH PACIFIC SEA LEVEL & CLIMATE MONITORING PROJECT

MONTHLY DATA REPORT

NO. 193

JULY 2011

Australian Government

Bureau of Meteorology

This project is sponsored by the Australian Agency for International Development (AusAID), and is managed by the Bureau of Meteorology with its National Tidal Centre (NTC) providing key technical support.

Australian Government

Bureau of Meteorology

National Tidal Centre
Bureau of Meteorology
Australia

GPO Box 421
Kent Town SA 5071
Australia

Tel: (+618) 8366 2730
Fax: (+618) 8366 2651
Website: http://www.bom.gov.au/oceanography/

Quality Certification:

I authorise the issue of this South Pacific Sea Level and Climate Monitoring Project Monthly Data Report for July 2011 in accordance with National Tidal Centre Quality Assurance procedures.

William Mitchell
Manager - National Tidal Centre

South Pacific Sea Level and Climate Monitoring Project

Monthly Data Report

July 2011

EXECUTIVE SUMMARY

This summary, and the overview that follows, are intended to provide a synopsis of the Monthly Data Report and of the trends observed over the life of the project to date.

July 2011

- The SEAFRAME network continued to collect high quality sea level and associated meteorological information for monitoring climate variability and climate change.

- El Niño - Southern Oscillation climate indicators were neutral across the equatorial Pacific during July. Subsurface ocean temperatures cooled slightly and periods of stronger than normal Trade Winds were observed, but overall conditions were near normal for this time of the year.

- Monthly sea levels for July 2011 were 5cm higher than normal at Marshall Islands, Samoa, Vanuatu, Fiji and Cook Islands and close to normal at the remaining SEAFRAME stations.

- International climate models predict that neutral climate conditions will persist through the southern hemisphere spring.

Short-Term Trends

It is important to stress that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Observed trends in sea level include natural variability, for example, events such as El Niño and effects due to many other atmospheric, oceanographic and geological processes. Longer-term data sets for all stations are required in order to separate the effects of the different signals. ***Please exercise caution in interpreting the short-term trends in the table below –*** they will almost certainly change over the coming years as the data set increases in length. Figure 13 later in this report provides the "time history" of the short-term trend at all project locations.

Recent short-term sea level trends in the project area based upon SEAFRAME data through July, 2011

Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cook Is	21°12'17.1"S / 159°47'5.2"W	Feb 1993	+4.9	+0.1
Tonga	21°8'12.5"S / 175°10'50.5"W	Jan 1993	+8.4	0.0
Fiji	17°36'17.7"S / 177°26'17.7"E	Oct 1992	+5.0	+0.1
Vanuatu	17°45'19.2"S / 168°18'27.7"E	Jan 1993	+5.3	+0.1
Samoa	13°49'36.4"S / 171°45'40.7"W	Feb 1993	+6.4	+0.1
Tuvalu	8°30'8.9"S / 179°11'42.6"E	Mar 1993	+4.0	0.0
Kiribati	1°21'54.2"N / 172°55'58.8"E	Dec 1992	+3.1	0.0
Nauru	0°31'45.9"S / 166°54'36.2"E	Jul 1993	+3.8	0.0
Solomon Is.	9°25'44.1"S / 159°57'19.3"E	Jul 1994	+7.1	0.0
PNG	2°2'31.5"S / 147°22'25.6"E	Sep 1994	+7.7	0.0
FSM	6°58'49.9"N / 158°12'0.8"E	Dec 2001	+17.0	+0.1
Marshall Is.	7°6'21.7"N / 171°22'22.1"E	May 1993	+4.8	+0.1

INTRODUCTION

Welcome to the July 2011 Monthly Data Report for the South Pacific Sea Level and Climate Monitoring Project (SPSLCMP). The report details the month by month operation of the SEAFRAME monitoring stations in the Pacific, including operational problems with the network or with satellite communications, the occurrence of abnormal sea level or climate events, interpretation of sea level fluctuations in the context of El Niño and the emergence of trends in the data.

The SPSLCMP was developed as an Australian response to concerns raised by the member countries of the South Pacific Forum over the potential impacts of global warming on climate and sea levels in the Pacific. Support was provided for the installation of SEAFRAME monitoring stations across the South Pacific Forum region.

SEAFRAME gauges not only measure sea level by two independent means, but also observe a number of "ancillary" variables - air and water temperatures, wind speed,

wind direction and atmospheric pressure. There is an associated programme of levelling to first order, to determine shifts in the vertical of the sea level sensors due to local land movement. Continuous Global Positioning System (CGPS) measurements are now also being made to determine the vertical movement of the land with respect to the International Terrestrial Reference Frame.

The AusAID funded project has, as its principal objective ***'the provision of an accurate long term record of sea level in the South Pacific for partner countries and the international scientific community, which enables them to respond to and manage, related impacts'***.

The project's monitoring network consists of 12 SEAFRAME stations, providing a wide coverage across the Southwest Pacific basin. All of these stations (see Figure A), with the exception of the Pohnpei (FSM) gauge, which was established in December 2001, have been operational since October 1994.

The monthly data report, one of a range of information products produced by the project, is the primary form of SPSLCMP data dissemination. Its content is designed to provide up-to-date access to the project's data products.



Figure A: *South Pacific Sea Level and Climate Monitoring Stations*

JULY CLIMATOLOGY

El Niño – Southern Oscillation (ENSO) climate indicators remained neutral across the equatorial Pacific during July. Subsurface ocean temperatures cooled slightly across the equatorial Pacific, but remain near normal for this time of the year. The Trade Winds and cloudiness over the central equatorial Pacific fluctuated around normal levels. International climate models predict neutral climate conditions will persist through the southern hemisphere winter and into spring.

The Southern Oscillation Index (SOI) was indicative of neutral climate conditions in June at +0.2, but rose in July to +10.7 (**Figure B**). Sustained positive values of the SOI above +8 are typical of La Niña, while sustained negative values below –8 are typical of El Niño.

Sea surface temperatures continued to warm slightly across the equatorial Pacific. Sea surface temperature anomalies during July were close to normal for this time of the year (**Figure C**). Slightly warmer than normal sea surface temperatures were observed in some areas of the southwest Pacific.

Subsurface ocean temperatures cooled slightly across the equatorial Pacific during July 2011. Cool subsurface anomalies appeared across parts of the central and eastern equatorial Pacific and the extent of warm subsurface anomalies decreased across the western equatorial Pacific (**Figure D**).

During El Niño (warm-episode) conditions there is a sustained weakening of the Trade Winds across much of the equatorial Pacific and an increase in cloudiness in the central equatorial Pacific particularly near the dateline. During La Niña (cold-episode) conditions there is a reversal of this situation, with stronger Trade Winds and a decrease in cloudiness in the central Pacific. Stronger than normal Trade Winds were observed across the equatorial Pacific during parts of July but were mostly close to normal for this time of the year (**Figure E**). Cloudiness near the dateline was also near neutral during July.

The consensus among international computer models surveyed by the Bureau of Meteorology suggests that the neutral climate conditions will persist through the southern hemisphere winter and into spring.

The preceding description of the climatology of the Pacific region, and Figures B, C and D are based on information sourced from the National Climate Centre of the Australian Bureau of Meteorology at http://www.bom.gov.au/climate/. Figure E was generated from the Tropical Atmosphere Ocean project website courtesy of PMEL, NOAA at http://www.pmel.noaa.gov/tao/.



Figure B: *The five-month weighted mean and individual monthly means of the Southern Oscillation Index (SOI). The SOI is ten times the monthly anomaly of the difference in mean sea level pressure between Tahiti and Darwin, divided by the long-term standard deviation of that difference for the relevant month.*



Figure C: *Sea surface temperature anomaly (°C) for July 2011.*



Figure D: *Equatorial depth-longitude section of ocean temperature anomalies for April through to July 2011. Contour interval is 0.5°C.*



Figure E: *Monthly mean wind vectors (top) and anomalies (bottom) for July 2011. The colour-shaded contours represent the monthly mean sea surface temperatures (top) and anomalies (bottom).*

JULY SEAFRAME DATA

Monthly Sea Level and Environmental Data (Figures 1-10)

The **observed sea levels (Figure 1)** are dominated by the daily oscillations of the tide. In most cases, the tide rises and falls twice per day (semi-diurnal), but at PNG and the Solomon Islands the tide tends to have a single high and low per day (diurnal). Where the tides are semi-diurnal the greatest tidal variations are called spring tides, which tend to occur close to the full and new moon. There was a new moon on the 1st of July and a full moon on the 15th of July, which was followed by another new moon on the 30th of July UTC.

Gaps in the data are the result of instrumental errors or data retrieval problems and are discussed under **Instrument Performance**.

The **residuals (Figure 2)** are the differences between the observed sea levels and the tidal predictions. They highlight non-tidal sea level fluctuations, such as those due to the effects of weather or tsunamis. Tropical cyclones often produce storm surges where the combination of low barometric pressure and strong winds raise sea levels well above the predicted tides for a period of a day or more.

The non-tidal sea level fluctuations can be amplified or sustained by the shape of the harbour in which the gauge is located. Some of the SEAFRAME stations are located in harbours that are favourable to persistent 'sloshing' under certain conditions (a phenomenon referred to as a seiche), such as at PNG when the wind suddenly changes strength or direction, at FSM during periods of reduced tidal range and at Nauru during strong westerly winds.

The sea level residuals at all stations, to some degree, exhibit semi-diurnal or diurnal fluctuations, which last a few days or weeks and then disappear. If these peaks were to persist, rather than appear as occasional 'transients', then the tidal analysis would be able to account for them, and the end result would be virtual eradication from the residuals.

The **barometrically corrected residuals (Figure 3)** have had the effect of atmospheric pressure fluctuations removed from the sea level residuals of Figure 2. The rule of thumb for the 'inverse barometer effect' is that a 1-hPa fall in the barometer, if sustained over a day or more, produces a 1-cm rise in the local sea level (within the area beneath the low pressure system).

The **winds**, **temperatures** and **barometric pressures** are plotted in **Figures 4** to **9**. The short lines in **Figure 5** follow the meteorological convention, that is, they point in the direction the wind is coming *from*. For example, the winds at Marshall Islands prevailed from the northeast for most of the month.

Air and water temperatures **(Figures 7** and **8)** are plotted using the same vertical scale for the purpose of comparison. The air temperatures are seen to fluctuate over a much wider range than the water temperatures. At some sites (e.g. FSM) the water temperature shows almost no variation, although the air temperature varies by several degrees between night and day. At Nauru a twice-daily fluctuation in water temperature is related to interactions between tides and terrestrial (land-based) water

discharging into the wharf area. The water temperature fluctuations there are usually more pronounced during the larger spring tides.

Barometric pressures **(Figure 9)** tend to fluctuate by around 3 hPa twice-daily at all stations as a result of atmospheric tides, which are largest in the tropical regions and reduce to near zero toward the poles. The longer-term barometric pressure fluctuations that occur over periods of days to weeks are due to passing weather systems. These fluctuations tend to be larger at sites farther away from the equator such as Cook Islands and Tonga.

The **meteorological data** are put into perspective by **Figure 10**. In this figure, if an open circle falls above (below) a solid dot, a new maximum (minimum) for the particular month has been set. *The data sets only include South Pacific Sea Level and Climate Monitoring Project data, which have been collected since October 1992 when the first station was installed (Fiji). The data from FSM has only been collected since December 2001.* Record July maximum air temperatures were observed at Samoa (32.1ºC), Fiji (29.7ºC) and Tonga (27.0ºC). Record July minimum barometric pressures were observed at Marshall Islands (1003.4 hPa), Solomon Islands (1004.6 hPa), Kiribati (1002.5 hPa), Tuvalu (1004.1 hPa) and Vanuatu (1002.5 hPa).

Mean Sea Level and Anomalies (Figures 11-13)

Figure 11 shows the **monthly mean sea levels**, which are simple arithmetic averages of the sea levels, relative to an arbitrary zero. The figure shows that Tuvalu, for example, normally experiences an annual cycle of about 0.2 metres, reaching a peak around February or March. One effect of the El Niño of 1997/1998 was very low sea levels which disrupted the annual sea level cycle at many of the SEAFRAME stations.

Figure 12 shows the monthly mean **sea level anomalies,** or departures from normal conditions after tides, annual and semi-annual seasonal cycles and the sea level trend have been removed. The annual cycle at Tuvalu (which has the largest consistent annual cycle) is quite notable in **Figure 11** but less apparent in **Figure 12**. By removing the seasonal cycles, the anomalies help to bring out irregular features, such as lower than normal sea levels across the region during the 1997/98 El Niño.

Monthly mean sea levels during July 2011 were around 5cm higher than normal at Marshall Islands, Samoa, Vanuatu, Fiji and Cook Islands. Sea levels recorded at other SEAFRAME stations during July were close to normal for this time of the year.

Sea Level Trends

The **short-term sea level trends** at individual stations as at July 2011 are shown in the following table. Sea level trends are updated every month by allowing for a linear trend term in the tidal analysis of all the data available at individual stations. *Please exercise caution in interpreting the trends* – they will continue to change over the coming years as the data sets increase in length. The evolution of the monthly trend values (in mm per year) at each station from one year after installation to present is depicted in **Figure 13**. This figure illustrates that as the sea level record becomes longer, the relative sea level trend estimates become more stable and reliable. The reason for this is that the trends from short sea level records are affected by the natural sea level variability occurring on inter-annual, El Niño and decadal timescales due to atmospheric, oceanographic and geological processes. Longer-term data sets for all

stations are required in order for the underlying trend to emerge from these short-term variations. Further details are available from the *National Tidal Centre (NTC), Australian Bureau of Meteorology*.

Recent short-term sea level trends in the project area based upon SEAFRAME data through July, 2011

Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cook Is	**21°12'17.1"S / 159°47'5.2"W**	**Feb 1993**	**+4.9**	**+0.1**
Tonga	**21°8'12.5"S / 175°10'50.5"W**	**Jan 1993**	**+8.4**	**0.0**
Fiji	**17°36'17.7"S / 177°26'17.7"E**	**Oct 1992**	**+5.0**	**+0.1**
Vanuatu	**17°45'19.2"S / 168°18'27.7"E**	**Jan 1993**	**+5.3**	**+0.1**
Samoa	**13°49'36.4"S / 171°45'40.7"W**	**Feb 1993**	**+6.4**	**+0.1**
Tuvalu	**8°30'8.9"S / 179°11'42.6"E**	**Mar 1993**	**+4.0**	**0.0**
Kiribati	**1°21'54.2"N / 172°55'58.8"E**	**Dec 1992**	**+3.1**	**0.0**
Nauru	**0°31'45.9"S / 166°54'36.2"E**	**Jul 1993**	**+3.8**	**0.0**
Solomon Is.	**9°25'44.1"S / 159°57'19.3"E**	**Jul 1994**	**+7.1**	**0.0**
PNG	**2°2'31.5"S / 147°22'25.6"E**	**Sep 1994**	**+7.7**	**0.0**
FSM	**6°58'49.9"N / 158°12'0.8"E**	**Dec 2001**	**+17.0**	**+0.1**
Marshall Is.	**7°6'21.7"N / 171°22'22.1"E**	**May 1993**	**+4.8**	**+0.1**

Barometric Pressure, Water Temperature and Air Temperature Anomalies

The anomalies of barometric pressure, water and air temperature (**Figures 14** to **16**) are determined in the same manner as the sea level anomalies (**Figure 12**), except the trend is not calculated.

The **barometric pressure anomalies (Figure 14)** show substantially higher than normal barometric pressures were observed at SEAFRAME stations during the 1997-1998 El Niño. July 2011 barometric pressures were generally close to normal for this time of the year. At Marshall Islands, PNG, Kiribati, Nauru, Vanuatu and Tonga monthly barometric pressures were slightly lower than normal but within -1 hPa of normal. At Cook Islands, the monthly barometric pressure was around +1 hPa higher than normal.

Water temperature anomalies (Figure 15) were near +1ºC at Vanuatu, Fiji, Tonga and Cook Islands during July 2011, indicating warmer than normal conditions. Water temperatures at the other SEAFRAME stations were generally close to normal for this time of the year.

Air temperature anomalies (Figure 16) at Fiji, Tonga and Cook Islands during July 2011 were also around +1ºC, while at Samoa and Vanuatu they were closer to +0.5 ºC. Air temperatures were close to normal at the other stations. Over the duration of the record the air temperature anomalies generally (although not always) follow the water temperature anomalies, which is an indication of the large influence the ocean has upon the climate of the Pacific Islands.

Instrument Performance

In **Figure 17,** which shows **sea level data return,** colour is used to distinguish five-year project phases. The number of missing days is noted in gaps in the bars.

Sea level data return during July 2011 was very high for all stations. Calibration and maintenance servicing was performed at Marshall Islands. Data that had not been transmitted from Tonga from 11th May through to 14th June 2011 has now been recovered from the data logger memory card. Minor satellite and dial-up communications problems resulted in small data loss from FSM, Nauru, Tuvalu and Samoa. Problems encountered with the ancillary sensors during July included the air temperature sensors at Nauru and Tuvalu, the water temperature sensors at Tuvalu and Tonga and the wind monitors at Kiribati, Vanuatu and Tuvalu. The erroneous data received from these problematic sensors were removed from the archived records.

Web Hits

The following chart shows the number of times the Pacific pages on the *NTC* web site have been visited, by month since January 2008.



SEAFRAME STATIONS

SEAFRAME stations employ either a SUTRON or TELMET (for upgraded stations) programmable data logger, water level gauges and other sensors. The data logger and associated electronics are normally housed in fibreglass huts. A sketch of a typical SUTRON station is shown in the following figure.

Water level sensors include:

(1) Primary water level using a Bartex 'AQUATRAK' acoustic-in-air sensor,

(2) Secondary water level (or backup) using a Druck pressure transducer mounted close to the seabed, and

(3) Tertiary water level using a Vega-puls radar sensor mounted above the water (at upgraded sites).

For SUTRON stations, the water level samples are averaged over three minutes and logged every six minutes, while meteorological sensors are logged on an hourly basis. With the upgraded TELMET stations, the water level samples are averaged over one minute and, together with meteorological data, logged every minute. Appropriate weighted-average and time-centred data is computed remotely which conforms to the SUTRON algorithm. Both SUTRON and TELMET data loggers have the memory capacity to store approximately one month of data.



The Observation Network Upgrade Project (ONUP) is scheduled to upgrade all Pacific SEAFRAME stations by mid-2013 with modernised TELMET data loggers, real-time satellite communications and additional radar-type water level sensors. The status of the station upgrades is given in the following table.

Status of Station Equipment Upgrades to July, 2011

Location	Lat / Long	SUTRON Installation Date	TELMET Upgrade Date
Cook Is	21°12'17.1"S / 159°47'5.2"W	Feb 1993	To be upgraded
Tonga	21°8'12.5"S / 175°10'50.5"W	Jan 1993	Mar 2011
Fiji	17°36'17.7"S / 177°26'17.7"E	Oct 1992	June 2011
Vanuatu	17°45'19.2"S / 168°18'27.7"E	Jan 1993	To be upgraded
Samoa	13°49'36.4"S / 171°45'40.7"W	Feb 1993	To be upgraded
Tuvalu	8°30'8.9"S / 179°11'42.6"E	Mar 1993	To be upgraded
Kiribati	1°21'54.2"N / 172°55'58.8"E	Dec 1992	To be upgraded
Nauru	0°31'45.9"S / 166°54'36.2"E	Jul 1993	To be upgraded
Solomon Is.	9°25'44.1"S / 159°57'19.3"E	Jul 1994	To be upgraded
PNG	2°2'31.5"S / 147°22'25.6"E	Sep 1994	To be upgraded
FSM	6°58'49.9"N / 158°12'0.8"E	Dec 2001	To be upgraded
Marshall Is.	7°6'21.7"N / 171°22'22.1"E	May 1993	To be upgraded

The *Monthly Data Report* is prepared by *NTC* for *AusAID*.

NTC would appreciate feedback from readers on the content and presentation of the *Monthly Data Report*.

Please spare a few moments to let us know your constructive opinion.

Further communication on the *Monthly Data Report* may be made to *NTC*. Anyone interested in a more detailed account of the project should contact:

National Tidal Centre
Bureau of Meteorology
PO Box 421
Kent Town SA 5067
Tel: (+618) (08) 8366 2600
Fax: (+618) (08) 8366 2693
Website: http://www.bom.gov.au/oceanography/tides.shtml

Or visit the project website at http://www.bom.gov.au/pacificsealevel

Please refer to: http://www.bom.gov.au/oceanography/projects/spslcmp/spslcmp.shtml
for details.

Please also note the following:

While care has been taken in the collection, analysis, and compilation of the data, it is supplied on the condition that neither the *Commonwealth of Australia* nor *NTC* shall be liable for any loss or injury whatsoever arising from the use of the data. Copyright for material contained in this document is held by the *Commonwealth of Australia*.

Figure 1

JULY 2011
SIX MINUTE WATER LEVEL OBSERVATIONS (m)



Figure 2

JULY 2011
SIX MINUTE RESIDUAL WATER LEVELS (m)



Figure 3

JULY 2011
SIX MINUTE RESIDUALS
ADJUSTED FOR ATMOSPHERIC PRESSURE (m)



Figure 4

JULY 2011
HOURLY WIND SPEEDS (m/s)



Figure 5

JULY 2011
HOURLY INCIDENT WINDS (m/s, deg True)

—— 10 m/s



Figure 6

JULY 2011
HOURLY MAXIMUM WIND GUSTS (m/s)



Figure 7

JULY 2011
HOURLY AIR TEMPERATURES (˚C)



Figure 8

JULY 2011
HOURLY WATER TEMPERATURES (°C)



Figure 9

JULY 2011
HOURLY ATMOSPHERIC PRESSURE (hPa)



Figure 10

Comparison of July 2011 Max, Min & Mean with Long Term July Values







Stations

1 - Marshall Islands
2 - Federated States of Micronesia
3 - Papua New Guinea
4 - Solomon Islands
5 - Kiribati
6 - Nauru
7 - Tuvalu
8 - Samoa
9 - Vanuatu
10 - Fiji
11 - Tonga
12 - Cook Islands

July 2011 Maximum
July 2011 Mean
July 2011 Minimum

Long Term July Maximum
Long Term July Mean
Long Term July Minimum

Figure 11

MONTHLY MEAN SEA LEVELS TO JULY 2011 (m)

The zero line represents an arbitrary fixed offset from the zero of the tide gauge.



Figure 12

SEA LEVEL ANOMALIES THROUGH JULY 2011 (m)



Figure 13

SEA LEVEL TRENDS THROUGH JULY 2011 (mm/year)



Figure 14

BAROMETRIC PRESSURE ANOMALIES THROUGH JULY 2011 (hPa)



Figure 15

WATER TEMPERATURE ANOMALIES THROUGH JULY 2011 (˚C)



Figure 16

AIR TEMPERATURE ANOMALIES THROUGH JULY 2011 (˚C)



Figure 17

SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF GAP ARE INDICATED

GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE

* Patchy record