

THE AUSTRALIAN BASELINE SEA LEVEL MONITORING PROJECT

MONTHLY DATA REPORT

MAY 2002



This report was prepared under the Australian Greenhouse Science Program for the Australian Greenhouse Office, supported by NTF Australia at the Flinders University of South Australia.



**National Tidal Facility Australia
Flinders University
Adelaide • Australia**

GPO Box 2100
Adelaide SA 5001
Australia

Tel: (+618) 8201 7532
Fax: (+618) 8201 7523
Email: ntf@flinders.edu.au
Website: <http://www.ntf.flinders.edu.au>

Quality Certification:

I authorise the issue of this Australian Baseline Sea Level Monitoring Project Monthly Data Report for May 2002 in accordance with National Tidal Facility Australia Quality Assurance procedures.

Wolfgang Scherer
Director - National Tidal Facility Australia

The Australian Baseline Sea Level Monitoring Project

Monthly Data Report

MAY 2002

NOTES ON THE DATA FOR MAY 2002

Sea level data return this month was excellent at all stations.

The air temperature sensor at Thevenard remained faulty and the data has consequently been removed from the record.

The wind speed and wind gusts at Thevenard and Stony Point were recording zero values for relatively long periods throughout May. As a consequence these erroneous values have been removed (Figures 4 and 6). Calibration and maintenance visits to Cocos Islands, Esperance and Hillarys were completed in May.

The sea level anomalies (Figure 10) were negative at all sites except for Cocos Islands. The barometric pressure anomalies (Figure 11) were negative for most sites during May with the exception of Groote Eylandt, Darwin, Burnie and Spring Bay, which returned slightly positive values.

The residuals (Figures 2 and 3), or difference between the observations and the tidal predictions, are the non-tidal components of the sea level observations. The residuals are primarily the consequence of short-term meteorological effects (Figures 5 and 9) and may give the result of elevated or depressed sea level observations.

With regard to the water and air temperature anomalies in Figures 12 and 13 respectively, it must be noted that there are large gaps in the data for several stations, where the data collected appeared to be erroneous. Also note that for several stations there were no backup water temperature sensors in operation, so the quality of this data is unknown. Similarly, air temperatures are compared to the temperature recorded by a sensor located in the upper levels of the environmental housing of the tide gauge. These will not exactly agree, as in locations where the housing is in the sun, the housing temperature will be higher than the actual air temperature. The temperature fluctuations inside the housing will also be less pronounced compared to the actual temperature fluctuations. This is due to the smaller amount of ventilation within the environmental housing. This can be used as a rough guide in determining the quality of air temperature data, but it is not an exact measure, and so is not used to fill the gaps.

It is difficult to relate the water and air temperature anomalies directly to those of barometric pressure and sea level without considering other effects, such as localised currents, wind speeds and directions. However, the anomalies are very useful in controlling the quality of the water and air temperature data at the Baseline stations.

Figure 16 compares the mean, maximum and minimum values for air temperature, water temperature and barometric pressure for the current month with the long-term May values. Please remember that the long-term ranges are calculated using the historical sets of May data for each station *excluding* the current month of data.

For most stations the mean air temperatures for May 2002 were consistent with the long term May means. Record maximum temperatures however were recorded at Cocos Islands, Esperance, Pt. Stanvac, Portland and Cape Ferguson, while no record minimums were reported this month. Failure of the air temperature sensor at Thevenard meant that no comparison could be made.

The water temperatures recorded at all sites for May 2002 were quite consistent with the long-term May values. A record maximum of 20°C was recorded at Thevenard this month but no record minimums were reported.

The barometric pressures were consistent with the long-term May values at all sites for May 2002, with the exception of Darwin and Broome where record maximums and minimums were reported and Port Kembla where a record minimum barometric pressure was reported.


Figure 14 shows the short-term sea level trends for each SEAFRAME location included in the Australian Baseline Sea Level Monitoring Project. Table 1 lists the commencement of operation, the sea level trend for the entire record (plotted in Figure 14) and the change in trend with respect to the previous month's analysis.

Table 1: Tide gauge installation dates , short-term sea level trends and change in trend from the previous month for the Australian Baseline array to May 2002.

Location	Installation Date	Sea Level Trend (mm/yr)	Change from previous month
Cocos Islands	Sep 1992	+13.3	+0.5
Groote Eylandt	Sep 1993	+26.9	-1.2
Darwin	May 1990	+17.4	-0.4
Broome	Nov 1991	+22.2	-0.5
Hillarys	Nov 1991	+16.9	-0.3
Esperance	Mar 1992	+12.1	-0.1
Thevenard	Mar 1992	+10.0	-0.1
Port Stanvac	Jun 1992	+9.6	-0.1
Portland	Jul 1991	+5.3	-0.1
Lorne	Jan 1993	+4.3	-0.2
Stony Point	Jan 1993	+4.2	-0.2
Burnie	Sep 1992	+5.9	-0.1
Spring Bay	May 1991	+4.2	-0.1
Port Kembla	Jul 1991	+7.4	-0.1
Roslyn Bay	Jun 1992	+6.0	-0.3
Cape Ferguson	Sep 1991	+7.7	-0.2

Figure 17 shows the monthly mean sea levels with respect to an arbitrary fixed offset from the zero of the tide gauge. This plot clearly shows significant correlation in seasonal signals between stations, in contrast to the sea level anomalies plot, which has the seasonal signal removed from the data. Table 2 gives the number of hits to the Australian Baseline Sea Level Monitoring project web-pages from 1999 to May 2002.

Table 2: Number of hits on the Australian Baseline Sea Level Monitoring Project web pages from 1999 to 2002.



Please note: Tide gauges at Stony Point and Lorne do not record air temperature, water temperature and barometric pressure data and are not present in Figures 3,7,8,9,11,12,13 and 16. The tide gauge at Lorne does not record wind data and is not present in Figures 4,5 and 6.

The *Monthly Data Report* is prepared by NTF Australia for Environment Australia. Staff members produce the text, plots and tables.

Further information on the *Monthly Data Report* and other projects conducted by NTF Australia can be obtained from the following address.

Contact address: NTF Australia
 The Flinders University of South Australia
 GPO BOX 2100, Adelaide SA 5001
 Tel: [+61 8] 8201 7534
 Fax: [+61 8] 8201 7523
 Email: ntf@flinders.edu.au
 Website: <http://www.ntf.flinders.edu.au>

Please note the following:

While all care has been taken in the collection, analysis and compilation of the data, it is supplied on the condition that neither the *Commonwealth of Australia* nor *NTF Australia* shall be liable for any loss or injury whatsoever arising from the use of the data. The Commonwealth of Australia holds copyright for material contained in this document.

Individuals and organisations are advised that quality controlled six-minute or hourly data from these stations are available on request from NTF Australia. Some handling fees may be charged. For commercial agencies requesting data, some additional costs may be levied.

Figure 1
MAY 2002
SIX MINUTE OBSERVATIONS (m)




Figure 2
MAY 2002
SIX MINUTE RESIDUAL WATER LEVELS (m)




Figure 3
MAY 2002
SIX MINUTE RESIDUALS
ADJUSTED FOR ATMOSPHERIC PRESSURE (m)




Figure 4

MAY 2002
HOURLY WIND SPEEDS (m/s)




Figure 5

MAY 2002
HOURLY INCIDENT WINDS (m/s, deg True)




Figure 6

MAY 2002
HOURLY MAXIMUM WIND GUSTS (m/s)




Figure 7

MAY 2002

HOURLY AIR TEMPERATURES (°C)




Figure 8

MAY 2002

HOURLY WATER TEMPERATURES (°C)




Figure 9

MAY 2002
HOURLY ATMOSPHERIC PRESSURE (hPa)





Figure 10
SEA LEVEL ANOMALIES THROUGH MAY 2002 (m)




BAROMETRIC PRESSURE ANOMALIES THROUGH MAY 2002 (hPa)



WATER TEMPERATURE ANOMALIES THROUGH MAY 2002 (°C)



Figure 13
AIR TEMPERATURE ANOMALIES
THROUGH MAY 2002 (°C)



SEA LEVEL TRENDS THROUGH MAY 2002 (mm/year)



Figure 15 SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF MISSING DATA ARE INDICATED
GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE

* Patchy record





Figure 16
Comparison of May 2002 Max, Min & Mean with
Long Term May Values.



The zero line represents an arbitrary fixed offset from the zero of the tide gauge.

National Tidal Facility Australia