

**THE AUSTRALIAN BASELINE SEA LEVEL
MONITORING PROJECT**

MONTHLY DATA REPORT

FEBRUARY 2007



Australian Government

Bureau of Meteorology

This report was prepared under the Australian Greenhouse Science Program for the Australian Greenhouse Office, supported by the National Tidal Centre, Bureau of Meteorology.



Australian Government

Bureau of Meteorology

**National Tidal Centre
Bureau of Meteorology
Australia**

GPO Box 421
Kent Town, SA 5071
Australia

Tel: (+618) 8366 2730
Fax: (+618) 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Quality Certification:

I authorise the issue of this Australian Baseline Sea Level Monitoring Project Monthly Data Report for February 2007 in accordance with National Tidal Centre Quality Assurance procedures.

William Mitchell
Manager - National Tidal Centre

The Australian Baseline Sea Level Monitoring Project

Monthly Data Report

FEBRUARY 2007

INTRODUCTION

The mission of the Australian Baseline Sea Level Monitoring Project (ABSLMP) is to monitor changes in sea level around Australia. It involves the operation and maintenance of an array of high-resolution sea level gauges and associated meteorological instruments (see Figure B) and management of a quality controlled national database of observations that is made available to the scientific and wider communities.

The Baseline array and a similar array in the South Pacific have been widely acknowledged in the global science community as two of the most accurate and reliable sources for information on sea level and climate change anywhere in the world.

This report is one of a series of monthly data reports that provide tables and figures summarising the data collected to date. The accompanying text relates primarily to the quality of the data rather than its interpretation. Periodic scientific evaluation of the data in the context of climate variability and climate change is provided in an annual data report.

Trends are derived from the sea level record, however readers are cautioned against drawing any conclusions from short duration records, particularly when used in isolation from other phenomena. The sea level record includes natural variability, such as El Niño events and the effects of atmospheric, oceanographic and geological processes. It is important to note that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Vertical movement of the instrumentation relative to local topography is monitored and the results are listed on the Geoscience Australia web site. Movement of each station relative to the International Terrestrial Reference Frame is not monitored.

NOTES ON THE DATA FOR FEBRUARY 2007

Sea level data return (Figures 1 and 17) in February 2007 was good for most stations with the exception of Cape Ferguson where a gap remains in the data due to technical problems and Lorne where the tide gauge has been de-commissioned during wharf refurbishment.

The residuals (Figures 2 and 3), the difference between the observations and the tidal predictions, are the non-tidal components of the sea level observations. The residuals are primarily the consequence of short-term meteorological effects (Figures 5 and 9) and can also indicate the passage of a tsunami.

Figure 10 compares the mean, maximum and minimum values for air temperature, water temperature and barometric pressure for February 2007 with the long-term February values. Note that the long-term ranges are calculated using the previous sets of February data for each station *excluding* the current month of data.

The air and water temperatures in February 2007 were generally within the long-term February temperature ranges. The mean temperatures were however above the long term mean at most sites with the exception of Rosslyn Bay and Cape Ferguson where the mean temperatures were below the long-term means. Record maximum air temperatures were recorded at Groote Eylandt and Port Stanvac, a record minimum air temperature was also seen at Broome. Record maximum water temperatures were seen at Broome, Hillarys, Portland, Burnie and Spring bay. The barometric pressures in February 2007 were generally within the long-term February pressures.

Figure 11 shows the monthly mean sea levels with respect to an arbitrary fixed offset from the zero of the tide gauge. The monthly mean sea levels contain seasonal variations, in contrast to the sea level anomalies (Figure 12), which have the seasonal signals removed from the data.

The sea level anomalies (Figure 12) were near normal to negative from Hillarys in the South West around the northern seaboard to Port Kembla on the East coast. The anomalies changed from slightly positive to slightly negative at sites across the southern seaboard and at Cape Ferguson from slightly negative to slightly positive in February.

Figure 13 shows the history of the short-term sea level trend for each site during the life of the Australian Baseline Sea Level Monitoring Project. Table 1 lists the commencement of operation, the latest sea level trend, and the change in trend with respect to the previous month's analysis. The sea level trends decreased over the previous months trends at most sites with the exception of Cape Ferguson and Groote Eylandt where the trend increased by less than 0.1 mm/year over the January 2007 trend.

The barometric pressure anomalies (Figure 14) were near normal at most sites in February 2007 and positive at Portland, Burnie and Spring Bay.

It is difficult to relate the water and air temperature anomalies (Figures 15 and 16) directly to those of barometric pressure and sea level without considering other effects, such as localised currents, wind speeds and directions. The anomalies are primarily used to quality check the water and air temperature data.

The number of hits to the Australian Baseline Sea Level Monitoring project web pages from 2004 to February 2007 is given in Figure A.

Please note: Tide gauges at Stony Point and Lorne do not record air temperature, water temperature and barometric pressure data and are not present in Figures 3,7,8,9,11,12,13 and 16. The tide gauge at Lorne does not record wind data and is not present in Figures 4, 5 and 6.

Table 1: Tide gauge position, installation date, short-term sea level trends and change in trend from the previous month for the Australian Baseline array through February 2007. Note that the trend for Lorne has not been calculated due to lack of current sea level data.

Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cocos Islands	12°07'00"S / 96°53'31"E	Sep 1992	+9.1	-0.3
Groote Eylandt	13°51'36"S / 136°24'57"E	Sep 1993	+5.7	0.0
Darwin	12°28'19"S / 130°50'45"E	May 1990	+6.7	-0.1
Broome	18°00'03"S / 122°13'06"E	Nov 1991	+8.5	-0.2
Hillarys	31°49'32"S / 115°44'19"E	Nov 1991	+7.3	-0.1
Esperance	33°52'24"S / 121°53'42"E	Mar 1992	+4.5	-0.1
Thevenard	32°09'02"S / 133°38'25"E	Mar 1992	+3.0	-0.1
Port Stanvac	35°06'35"S / 138°27'55"E	Jun 1992	+4.2	-0.2
Portland	38°20'38"S / 141°36'49"E	Jul 1991	+2.1	-0.1
Lorne	38°30'00"S / 143°59'00"E	Jan 1993	na	na
Stony Point	38°22'00"S / 145°13'00"E	Jan 1993	+1.0	-0.2
Burnie	41°03'00"S / 145°54'53"E	Sep 1992	+1.9	-0.1
Spring Bay	42°32'47"S / 147°55'51"E	May 1991	+2.9	-0.2
Port Kembla	34°28'26"S / 150°54'43"E	Jul 1991	+3.4	-0.1
Rossllyn Bay	23°09'40"S / 150°47'24"E	Jun 1992	+1.4	-0.1
Cape Ferguson	19°16'39"S / 147°03'31"E	Sep 1991	+2.7	+0.1

Figure A: Number of hits on the Australian Baseline Sea Level Monitoring Project web pages from 2004 to February 2007.





Figure B: Australian Baseline Sea Level Monitoring Project sites.

The *Monthly Data Report* is prepared by the NTC, Bureau of Meteorology for Environment Australia. Staff members produce the text, plots and tables.

Further information on the *Monthly Data Report* and other projects conducted by the NTC, Bureau of Meteorology can be obtained from the following address.

National Tidal Centre
Bureau of Meteorology
GPO BOX 421, Kent Town SA 5071
Tel: [+61 8] 8366 2730
Fax: [+61 8] 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Please note the following:

While all care has been taken in the collection, analysis and compilation of the data, it is supplied on the condition that neither the *Commonwealth of Australia* nor the NTC, Bureau of Meteorology shall be liable for any loss or injury whatsoever arising from the use of the data. The Commonwealth of Australia holds copyright for material contained in this document.

Individuals and organisations are advised that quality controlled six-minute or hourly data from these stations are available on request from the NTC, Bureau of Meteorology. Some handling fees may be charged. For commercial agencies requesting data, some additional costs may be levied.

Figure 1

FEBRUARY 2007
SIX MINUTE SEA LEVEL OBSERVATIONS (m)




Figure 2

FEBRUARY 2007
SIX MINUTE RESIDUAL WATER LEVELS (m)




Figure 3
FEBRUARY 2007
SIX MINUTE RESIDUALS
ADJUSTED FOR ATMOSPHERIC PRESSURE (m)




Figure 4

**FEBRUARY 2007
HOURLY WIND SPEEDS (m/s)**




Figure 5

FEBRUARY 2007
HOURLY INCIDENT WINDS (m/s, deg True)





Figure 6

FEBRUARY 2007
HOURLY MAXIMUM WIND GUSTS (m/s)



February 2007 (UTC)

Figure 7

FEBRUARY 2007
HOURLY AIR TEMPERATURES (°C)




Figure 8

FEBRUARY 2007
HOURLY WATER TEMPERATURES (°C)




Figure 9

FEBRUARY 2007
HOURLY ATMOSPHERIC PRESSURE (hPa)







Figure 10
Comparison of February 2007 Max, Min & Mean with
Long Term February Values.



Stations

- 1 - Cocos Islands
- 2 - Groote Eylandt
- 3 - Darwin
- 4 - Broome
- 5 - Hillarys
- 6 - Esperance
- 7 - Thevenard
- 8 - Port Stanvac
- 9 - Portland
- 10 - Burnie
- 11 - Spring Bay
- 12 - Port Kembla
- 13 - Rosslyn Bay
- 14 - Cape Ferguson

- February 2007 Maximum
- February 2007 Mean
- February 2007 Minimum
- Long Term February Maximum
- Long Term February Mean
- Long Term February Minimum

Figure 11

MONTHLY MEAN SEA LEVELS TO FEBRUARY 2007 (m)

The zero line represents an arbitrary fixed offset from the zero of the tide gauge.




Figure 12
SEA LEVEL ANOMALIES THROUGH FEBRUARY 2007 (m)




Figure 13

SEA LEVEL TRENDS THROUGH FEBRUARY 2007 (mm/year)




Figure 14

BAROMETRIC PRESSURE ANOMALIES THROUGH FEBRUARY 2007 (hPa)




Figure 15

WATER TEMPERATURE ANOMALIES THROUGH FEBRUARY 2007 (°C)




Figure 16
AIR TEMPERATURE ANOMALIES
THROUGH FEBRUARY 2007 (°C)




Figure 17 SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF MISSING DATA ARE INDICATED
GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE

* Patchy record

