

**THE AUSTRALIAN BASELINE SEA LEVEL
MONITORING PROJECT**

MONTHLY DATA REPORT

AUGUST 2008



Australian Government

Bureau of Meteorology

This report was prepared under the Australian Greenhouse Science Program for the Australian Greenhouse Office, supported by the National Tidal Centre, Bureau of Meteorology.



Australian Government

Bureau of Meteorology

**National Tidal Centre
Bureau of Meteorology
Australia**

GPO Box 421
Kent Town, SA 5071
Australia

Tel: (+618) 8366 2730
Fax: (+618) 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Quality Certification:

I authorise the issue of this Australian Baseline Sea Level Monitoring Project Monthly Data Report for August 2008 in accordance with National Tidal Centre Quality Assurance procedures.

William Mitchell
Manager - National Tidal Centre

The Australian Baseline Sea Level Monitoring Project

Monthly Data Report

AUGUST 2008

INTRODUCTION

The mission of the Australian Baseline Sea Level Monitoring Project (ABSLMP) is to monitor changes in sea level around Australia. It involves the operation and maintenance of an array of high-resolution sea level gauges and associated meteorological instruments (see Figure B) and management of a quality controlled national database of observations that is made available to the scientific and wider communities.

The Baseline array and a similar array in the South Pacific have been widely acknowledged in the global science community as two of the most accurate and reliable sources for information on sea level and climate change anywhere in the world.

This report is one of a series of monthly data reports that provide tables and figures summarising the data collected to date. The accompanying text relates primarily to the quality of the data rather than its interpretation. Periodic scientific evaluation of the data in the context of climate variability and climate change is provided in an annual data report.

The trends are derived from the sea level record. However, readers are cautioned against drawing any conclusions from short duration records, particularly when used in isolation from other phenomena. The sea level record includes natural variability, such as El Niño events and the effects of atmospheric, oceanographic and geological processes. It is important to note that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Vertical movement of the instrumentation relative to local topography is monitored and the results are listed on the Geoscience Australia web site. Movement of each station relative to the International Terrestrial Reference Frame is not monitored.

NOTES ON THE DATA FOR AUGUST 2008

Sea level data return (Figures 1 and 17) in August 2008 was good for most stations. Due to occupational health and safety reasons the station at Broome is being switched off when ships are being loaded. This has resulted in a loss of three days of un-recoverable data.

The residuals (Figures 2 and 3), the difference between the observations and the tidal predictions, are the non-tidal components of the sea level observations. The residuals are primarily the consequence of short-term meteorological effects (Figures 5 and 9) and can also indicate the passage of a tsunami. The meteorological convention is followed in Figure 5 where the vector indicates the direction from which the wind is coming.

Figure 10 compares the mean, maximum and minimum values for air temperature, water temperature and barometric pressure for August 2008 with the long-term values. Note that

the long-term ranges are calculated using the previous sets of August data for each station **excluding** the current month of data.

The mean air temperatures in August 2008 were slightly lower than the long term August mean. A record low monthly air temperature of 3.9°C was set at Port Stanvac this month. Generally, the mean water temperatures for August 2008 were lower than the long-term mean but no site exceeded the upper or lower long-term August limits. The mean barometric pressures for August 2008 were greater than the long-term means for all sites except Cocos Islands. The minimum barometric pressures, anti-clockwise around Australia from Hillarys to Cape Ferguson were substantially higher than the long-term minima, with record maximum August barometric pressures recorded at Hillarys (1037.1 hPa) and Esperance (1040.8 hPa). These values compare with Bureau of Meteorology observations at Perth (1038.0hPa) and Esperance (1040.1hPa) recorded on the 21st of August.

Figure 11 shows the monthly mean sea levels with respect to an arbitrary fixed offset from the zero of the tide gauge. The monthly mean sea levels contain seasonal variations, in contrast to the sea level anomalies (Figure 12), which have the seasonal signals removed from the data.

The sea level anomalies (Figure 12) in August 2008 were positive at Cocos Islands and Burnie, and slightly neutral to negative at the remaining project sites.

Figure 13 shows the history of the short-term sea level trend for each site during the life of the Australian Baseline Sea Level Monitoring Project. Table 1 lists the commencement of operation, the latest sea level trend and the change in trend with respect to the previous month's analysis. It is important to stress that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Observed trends in sea level include natural variability, for example, events such as El Niño and effects due to many other atmospheric, oceanographic and geological processes. Longer-term data sets for all stations are required in order to separate the effects of the different signals. ***Please exercise caution in interpreting the short-term trends in the table below*** – they will almost certainly change over the coming years as the data set increases in length.

The barometric pressure anomalies (Figure 14) in August 2008 reflect the positive displacement of the barometric pressures presented in Figure 10. Positive anomalies were recorded from Broome to Cape Ferguson with neutral anomalies recorded at Cocos Islands, Groote Eylandt and Darwin. It is difficult to relate the water and air temperature anomalies (Figures 15 and 16) directly to those of barometric pressure and sea level without considering other effects, such as localised currents, wind speeds and directions. The anomalies are primarily used to quality check the water and air temperature data.

The number of hits to the Australian Baseline Sea Level Monitoring project web pages from 2005 to August 2008 are given in Figure A.

Please note: Tide gauges at Stony Point and Lorne do not record air temperature, water temperature and barometric pressure data and are not present in Figures 3,7,8,9,11,12,13 and 16. The tide gauge at Lorne does not record wind data and is not present in Figures 4, 5 and 6.

Table 1: Tide gauge position, data start date, short-term sea level trends and change in trend from the previous month for the Australian Baseline array through August 2008.

Location	Lat / Long	Start Date	Trend (mm/yr)	Change from previous month
Cocos Islands	12°07'0.1"S / 96°53'30.9"E	Sep 1992	+8.3	+0.1
Groote Eylandt	13°51'36.2"S / 136°24'56.1"E	Sep 1993	+6.2	-0.1
Darwin	12°28'18.4"S / 130°50'45.1"E	May 1990	+7.2	0.0
Broome	18°00'03.0"S / 122°13'07.1"E	Nov 1991	+8.6	0.0
Hillarys	31°49'32.0"S / 115°44'18.9"E	Nov 1991	+8.5	0.0
Esperance	33°52'15.2"S / 121°53'43.3"E	Mar 1992	+5.4	-0.1
Thevenard	32°08'56.2"S / 133°38'28.8"E	Mar 1992	+3.7	-0.1
Port Stanvac	35°06'31.0"S / 138°28'1.3"E	Jun 1992	+4.9	-0.1
Portland	38°20'36.4"S / 141°36'47.4"E	Jul 1991	+2.6	-0.1
Lorne	38°32'49.4"S / 143°59'19.8"E	Jan 1993	+1.5	-0.1
Stony Point	38°22'19.7"S / 145°13'28.9"E	Jan 1993	+1.7	-0.1
Burnie	41°03'0.3"S / 145°54'54.0"E	Sep 1992	+2.0	0.0
Spring Bay	42°32'45.1"S / 147°55'57.8"E	May 1991	+3.2	-0.1
Port Kembla	34°28'25.5"S / 150°54'42.7"E	Jul 1991	+3.2	0.0
Roslyn Bay	23°09'39.7"S / 150°47'24.6"E	Jun 1992	+1.6	-0.1
Cape Ferguson	19°16'38.4"S / 147°03'30.4"E	Sep 1991	+2.7	-0.1

Figure A: Number of hits on the Australian Baseline Sea Level Monitoring Project web pages from 2004 to August 2008.





Figure B: Australian Baseline Sea Level Monitoring Project sites.



The *Monthly Data Report* is prepared by the NTC, Bureau of Meteorology for the Australian Greenhouse Office. Staff members produce the text, plots and tables.

Further information on the *Monthly Data Report* and other projects conducted by the NTC, Bureau of Meteorology can be obtained from the following address.

National Tidal Centre
Bureau of Meteorology
GPO BOX 421, Kent Town SA 5071
Tel: [+61 8] 8366 2730
Fax: [+61 8] 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Please note the following:

While all care has been taken in the collection, analysis and compilation of the data, it is supplied on the condition that neither the *Commonwealth of Australia* nor the NTC, Bureau of Meteorology shall be liable for any loss or injury whatsoever arising from the use of the data. The Commonwealth of Australia holds copyright for material contained in this document.

Individuals and organisations are advised that quality controlled six-minute or hourly data from these stations are available on request from the NTC, Bureau of Meteorology. Some handling fees may be charged. For commercial agencies requesting data, some additional costs may be levied.

Figure 1

**AUGUST 2008
SIX MINUTE SEA LEVEL OBSERVATIONS (m)**




Figure 2
AUGUST 2008
SIX MINUTE RESIDUAL WATER LEVELS (m)




Figure 3
AUGUST 2008
SIX MINUTE RESIDUALS
ADJUSTED FOR ATMOSPHERIC PRESSURE (m)




Figure 4

AUGUST 2008
HOURLY WIND SPEEDS (m/s)




Figure 5

AUGUST 2008
HOURLY INCIDENT WINDS (m/s, deg True)




Figure 6

AUGUST 2008
HOURLY MAXIMUM WIND GUSTS (m/s)




Figure 7

AUGUST 2008
HOURLY AIR TEMPERATURES (°C)




Figure 8

AUGUST 2008
HOURLY WATER TEMPERATURES (°C)




Figure 9

AUGUST 2008
HOURLY ATMOSPHERIC PRESSURE (hPa)







Figure 10
Comparison of August 2008 Max, Min & Mean with
Long Term August Values.



Stations

- 1 - Cocos Islands
- 2 - Groote Eylandt
- 3 - Darwin
- 4 - Broome
- 5 - Hillarys
- 6 - Esperance
- 7 - Thevenard
- 8 - Port Stanvac
- 9 - Portland
- 10 - Burnie
- 11 - Spring Bay
- 12 - Port Kembla
- 13 - Rosslyn Bay
- 14 - Cape Ferguson

- August 2008 Maximum
- August 2008 Mean
- August 2008 Minimum
- Long Term August Maximum
- Long Term August Mean
- Long Term August Minimum

Figure 11

MONTHLY MEAN SEA LEVELS TO AUGUST 2008 (m)

The zero line represents an arbitrary fixed offset from the zero of the tide gauge.




Figure 12
SEA LEVEL ANOMALIES THROUGH AUGUST 2008 (m)




Figure 13

SEA LEVEL TRENDS THROUGH AUGUST 2008 (mm/year)




Figure 14

BAROMETRIC PRESSURE ANOMALIES THROUGH AUGUST 2008 (hPa)




Figure 15

WATER TEMPERATURE ANOMALIES THROUGH AUGUST 2008 (°C)




Figure 16
AIR TEMPERATURE ANOMALIES
THROUGH AUGUST 2008 (°C)




Figure 17 SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF MISSING DATA ARE INDICATED
GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE

* Patchy record

