

**THE AUSTRALIAN BASELINE SEA LEVEL
MONITORING PROJECT**

MONTHLY DATA REPORT

JUNE 2010



Australian Government

Bureau of Meteorology

This report was prepared under the Australian Climate Change Science Program for the Department of Climate Change and Energy Efficiency, supported by the National Tidal Centre, Bureau of Meteorology.



Australian Government

Bureau of Meteorology

**National Tidal Centre
Bureau of Meteorology
Australia**

GPO Box 421
Kent Town, SA 5071
Australia

Tel: (+618) 8366 2730
Fax: (+618) 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Quality Certification:

I authorise the issue of this Australian Baseline Sea Level Monitoring Project Monthly Data Report for June 2010 in accordance with National Tidal Centre Quality Assurance procedures.

William Mitchell
Manager - National Tidal Centre

The Australian Baseline Sea Level Monitoring Project

Monthly Data Report

JUNE 2010

INTRODUCTION

The mission of the Australian Baseline Sea Level Monitoring Project (ABSLMP) is to monitor changes in sea level around Australia. It involves the operation and maintenance of an array of high-resolution sea level gauges and associated meteorological instruments (see Figure B) and management of a quality controlled national database of observations that is made available to the scientific and wider communities.

This report is one of a series of monthly data reports that provide tables and figures summarising the data collected to date. The accompanying text relates primarily to the quality of the data rather than its interpretation. Periodic scientific evaluation of the data in the context of climate variability and climate change is provided in an annual data report.

The trends are derived from the sea level record. However, readers are cautioned against drawing any conclusions from short duration records, particularly when used in isolation from other phenomena. The sea level record includes natural variability, such as El Niño events and the effects of atmospheric, oceanographic and geological processes. It is important to note that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Vertical movement of the instrumentation relative to local topography is monitored and the results are listed on the Geoscience Australia web site. Movement of each station relative to the International Terrestrial Reference Frame is not monitored.

NOTES ON THE DATA FOR JUNE 2010

Sea level data return (Figures 1 and 17) in June 2010 was excellent for most stations. The Baseline array network modernisation project has been completed with Telmet 320 loggers in operation at all locations except the privately owned Lorne and Stony Point stations. A calibration and maintenance visit was made to Cocos Islands from the 28th of June to the 3rd of July. Ten hours of data was lost at Broome as a result of the Broome Port Authority's policy of switching off the power when fuel ships are in dock. Problems with the recording of wind data at Broome since December have continued into June. Inspection of the wind monitor revealed evidence of water incursion into the terminal block, corrosion and broken wiring. The water temperature sensor failure at Broome continued into June. The Esperance water temperature sensor was erroneously recording a constant value (17.6°C) during June and as a consequence constant values recorded during May were also removed from the record. The Stony Point wind data has been removed while suspect high wind speeds and gusts are investigated further, whilst wind data received from Groote Eylandt suggests instrument failure and have been removed from the record.

The residuals (Figures 2 and 3), being the difference between the observations and the tidal predictions, are the non-tidal components of the sea level. They are primarily the consequence of short-term meteorological effects (Figures 5 and 9) and can also indicate the passage of a tsunami.

The meteorological convention is followed in Figure 5 where the vector indicates the direction from which the wind is blowing.

Figure 10 compares the mean, maximum and minimum values for air temperature, water temperature and barometric pressure for June 2010 with the long-term values. Note that the long-term ranges are calculated using the previous sets of June data for each station **excluding** the current month of data.

Record June minimum air temperatures were recorded at Hillarys (4.8°C) and Esperance (4.2°C) whilst Spring Bay recorded a new June maximum water temperature (14.3°C) and a new maximum June barometric pressure (1040.5hPa) this month.

Figure 11 shows the monthly mean sea levels with respect to an arbitrary fixed offset from the zero of the tide gauge. The monthly mean sea levels contain seasonal variations, in contrast to the sea level anomalies (Figure 12), which have the seasonal signals and trends removed from the data.

In June 2010 sea level anomalies (Figure 12) were close to zero or negative for all sites except Port Kembla, Rosslyn Bay and Cape Ferguson where the anomalies were greater than 5 cm.

Figure 13 shows the history of the short-term sea level trend for each site during the life of the Australian Baseline Sea Level Monitoring Project. Table 1 lists the commencement of operation, the latest sea level trend and the change in trend with respect to the previous month's analysis. It is important to stress that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Observed trends in sea level include natural variability, for example, events such as El Niño and effects due to many other atmospheric, oceanographic and geological processes. Longer-term data sets for all stations are required in order to separate the effects of the different signals. ***Please exercise caution in interpreting the short-term trends in the table below*** – they will almost certainly change over the coming years as the data set increases in length.

The barometric pressure anomalies (Figure 14) for June 2010 were positive along Southern Australia from Hillarys to Port Kembla and close to zero for all other locations. It is difficult to relate the water and air temperature anomalies (Figures 15 and 16) directly to those of barometric pressure and sea level without considering other effects, such as localised currents, wind speeds and directions. The anomalies are primarily used to quality check the water and air temperature data. Water temperature anomalies were close to zero for all locations except Thevenard, Port Stanvac, Burnie, Spring Bay and Port Kembla where the anomalies were strongly positive. Air temperature anomalies, which generally follow the water temperature anomalies, show close to normal temperatures were observed at all locations except Groote Eylandt or Broome where the anomalies were close to +1°C and Hillarys where the anomaly was around -1°C.

The number of hits to the Australian Baseline Sea Level Monitoring Project (ABSLMP) web pages from January 2007 to June 2010 are given in Figure A.

Please note: Tide gauges at Stony Point and Lorne do not record air temperature, water temperature and barometric pressure data and are not present in Figures 3,7,8,9,11,12,13 and 16. The tide gauge at Lorne does not record wind data and is not present in Figures 4, 5 and 6.

Table 1: Tide gauge position, data start date, short-term sea level trends and change in trend from the previous month for the Australian Baseline array through June 2010.

Recent short-term sea level trends in the project area based upon SEAFRAME data through June, 2010				
Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cocos Islands	12°07'07.1"S / 96°53'30.9"E	Sep1992	+8.7	-0.1
Groote Eylandt	13°51'36.2"S / 136°24'56.1"E	Sep 1993	+7.1	0.0
Darwin	12°28'18.4"S / 130°50'45.1"E	May 1990	+7.2	0.0
Broome	18°00'03.0"S / 122°13'07.1"E	Nov 1991	+7.8	0.0
Hillarys	31°49'32.0"S / 115°44'18.9"E	Nov 1991	+7.6	-0.1
Esperance	33°52'15.2"S / 121°53'43.3"E	Mar 1992	+5.3	-0.1
Thevenard	32°08'56.2"S / 133°38'28.8"E	Mar 1992	+4.2	-0.1
Port Stanvac	35°06'31.0"S / 138°28'1.3"E	Jun 1992	+5.0	-0.1
Portland	38°20'36.4"S / 141°36'47.4"E	Jul 1991	+3.0	0.0
Lorne	38°32'49.4"S / 143°59'19.8"E	Jan 1993	+1.3	0.0
Stony Point	38°22'19.7"S / 145°13'28.9"E	Jan 1993	+1.3	0.0
Burnie	41°03'0.3"S / 145°54'54.0"E	Sep 1992	+2.9	0.0
Spring Bay	42°32'45.1"S / 147°55'57.8"E	May 1991	+3.4	0.0
Port Kembla	34°28'25.5"S / 150°54'42.7"E	Jul 1991	+3.0	+0.1
Rosslyn Bay	23°09'39.7"S / 150°47'24.6"E	Jun 1992	+2.3	+0.1
Cape Ferguson	19°16'38.4"S / 147°03'30.4"E	Sep 1991	+3.4	+0.1

Figure A: Number of hits on the Australian Baseline Sea Level Monitoring Project web pages from 2007 to June 2010.





Figure B: Australian Baseline Sea Level Monitoring Project sites.



The *Monthly Data Report* is prepared by the NTC, Bureau of Meteorology for the Department of Climate Change and Energy Efficiency. Staff members produce the text, plots and tables.

Further information on the *Monthly Data Report* and other projects conducted by the NTC, Bureau of Meteorology can be obtained from the following address.

National Tidal Centre
Bureau of Meteorology
GPO BOX 421, Kent Town SA 5071
Tel: [+61 8] 8366 2730
Fax: [+61 8] 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Please note the following:

While all care has been taken in the collection, analysis and compilation of the data, it is supplied on the condition that neither the *Commonwealth of Australia* nor the NTC, Bureau of Meteorology shall be liable for any loss or injury whatsoever arising from the use of the data. The Commonwealth of Australia holds copyright for material contained in this document.

Individuals and organisations are advised that quality controlled six-minute or hourly data from these stations are available on request from the NTC, Bureau of Meteorology. Some handling fees may be charged. For commercial agencies requesting data, some additional costs may be levied.

Figure 1

JUNE 2010
SIX MINUTE SEA LEVEL OBSERVATIONS (m)




Figure 2
JUNE 2010
SIX MINUTE RESIDUAL WATER LEVELS (m)




Figure 3
JUNE 2010
SIX MINUTE RESIDUALS
ADJUSTED FOR ATMOSPHERIC PRESSURE (m)




Figure 4

JUNE 2010
HOURLY WIND SPEEDS (m/s)




Figure 5

JUNE 2010
HOURLY INCIDENT WINDS (m/s, deg True)




Figure 6

JUNE 2010
HOURLY MAXIMUM WIND GUSTS (m/s)




Figure 7

JUNE 2010
HOURLY AIR TEMPERATURES (°C)




Figure 8

JUNE 2010
HOURLY WATER TEMPERATURES (°C)




Figure 9

JUNE 2010
HOURLY ATMOSPHERIC PRESSURE (hPa)







Figure 10
Comparison of June 2010 Max, Min & Mean with
Long Term June Values.



Stations

- 1 - Cocos Islands
- 2 - Groote Eylandt
- 3 - Darwin
- 4 - Broome
- 5 - Hillarys
- 6 - Esperance
- 7 - Thevenard
- 8 - Port Stanvac
- 9 - Portland
- 10 - Burnie
- 11 - Spring Bay
- 12 - Port Kembla
- 13 - Rosslyn Bay
- 14 - Cape Ferguson

- June 2010 Maximum
- ◐ June 2010 Mean
- June 2010 Minimum
- Long Term June Maximum
- Long Term June Mean
- Long Term June Minimum

Figure 11

MONTHLY MEAN SEA LEVELS TO JUNE 2010 (m)

The zero line represents an arbitrary fixed offset from the zero of the tide gauge.




Figure 12
SEA LEVEL ANOMALIES THROUGH JUNE 2010 (m)




Figure 13

SEA LEVEL TRENDS THROUGH JUNE 2010 (mm/year)




Figure 14

BAROMETRIC PRESSURE ANOMALIES THROUGH JUNE 2010 (hPa)




Figure 15

WATER TEMPERATURE ANOMALIES THROUGH JUNE 2010 (°C)




Figure 16
AIR TEMPERATURE ANOMALIES
THROUGH JUNE 2010 (°C)




Figure 17 SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF MISSING DATA ARE INDICATED
GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE

* Patchy record

